

Neuropsychological impairment and brain damage in children and adolescents associated with preterm birth

Leire Zubiaurre-Elorza
Sara Soria
Carme Junqué
Universitat de Barcelona

Prematurity is defined by the World Health Organization (WHO, 1977) as a delivery occurring before 37 completed weeks of gestation. White matter (WM) abnormalities and periventricular leukomalacia (PVL) are the most prominent pathologies in the preterm neonate. Compared with their term-peers, children born preterm have lower intelligence quotient (IQ) scores, although within the normal range. Moreover, domain-specific measures of neuropsychological abilities such as executive functions, visuoperceptual and linguistic skills are often impaired. Preterm children with PVL have a similar profile, with lower IQ scores and poorer executive functions, mental calculation and visuoperceptual skills than controls. Cerebral structural gray matter (GM) abnormalities that might contribute to the poor performance of preterm samples are localized in the frontal, parietal, temporal and occipital lobes. The cerebellum and deep gray matter structures such as the caudate nuclei, the thalami and the hippocampus are also altered. Over the past few years, magnetic resonance imaging and neuropathological studies have shown that, in addition to the classical WM injury seen in preterm samples with PVL, cortical and subcortical GM injury may also contribute to the impaired cognitive performance.

Keywords: Intelligence, preterm, neuroimaging, neuropsychology, magnetic resonance imaging,

Alteraciones neuropsicológicas y daño cerebral en la infancia y adolescencia asociados al nacimiento prematuro

La prematuridad es definida por la Organización Mundial de la Salud (OMS, 1977) como el parto que ocurre antes de las 37 semanas de gestación. La alteración de la sustancia blanca (SB) es la principal característica de daño cerebral

y la leucomalacia periventricular (LPV) es el hallazgo patológico más común en estudios de resonancia magnética. Los niños nacidos prematuros comparados con los niños nacidos a término presentan un cociente de inteligencia (CI) con puntuaciones menores pero dentro de la normalidad. Las funciones neuropsicológicas específicas más afectadas son las ejecutivas, visuo-perceptivas y lingüísticas. Los niños prematuros con LPV, también muestran puntuaciones más bajas en el CI, en las funciones ejecutivas, en el cálculo mental y en las capacidades visuoperceptivas. Las alteraciones cerebrales estructurales de sustancia gris (SG) que pueden contribuir a una peor ejecución cognitiva en estas muestras de prematuros implican regiones en los lóbulos cerebrales temporal, parietal, frontal y occipital. Además, el cerebelo y diversos núcleos grises subcorticales tales como el núcleo caudado, el tálamo y el hipocampo están también afectados. En los últimos años, los estudios de resonancia magnética y neuropatológicos, están demostrando que además de la alteración clásica de SB, los niños prematuros con LPV presentan daño en la SG tanto cortical como subcortical a los cuales se les atribuye una implicación directa en la peor ejecución cognitiva.

Palabras clave: inteligencia, prematuro, neuroimagen, neuropsicología, resonancia magnética.

Introduction

The World Health Organization (World Health Organization [WHO], 1977) defines preterm birth as a delivery occurring prior to 37 completed weeks of gestation. The incidence of preterm birth in Spain has risen from 4.75% of all live births in 1996 to 6.88% in 2010 (National Statistics Institute of Spain, 2010, www.ine.es). The reasons for this increment in preterm deliveries remain unclear, although it has been proposed that an increase in multiple gestations associated with in vitro fertilization therapy is in part responsible (Blondel & Kaminski, 2002) as well as the improvements in antenatal and neonatal care (Simmons, Rubens, Darmstadt, & Gravett, 2010).

Preterm birth has negative consequences on normal brain maturation processes, which may lead to neurocognitive and behavioral problems in childhood (Bhutta, Cleves, Casey, Cradock, & Anand, 2002) and in adolescence and adulthood (Marlow *et al.*, 2004). The time window during childhood and adolescence is particularly sensitive; at this time age-related social and academic demands increase, and emerging sequelae manifest themselves or pre-existing ones are intensified (Aylward, 2005).

Neuropsychological profile in children/adolescents born preterm

A meta-analysis of studies published between 1980 and November 2001 (Bhutta *et al.*, 2002) focusing on the cognitive and behavioral outcomes of school-aged preterm children showed that preterm birth is associated with low cognitive scores in childhood. During the last decade, with the decrease in mortality and the

increasing recognition of neurodevelopmental disabilities, termed by Aylward (2002) as “high prevalence/low severity dysfunctions”, contemporary cohorts provide evidence of a similar pattern of low intelligence quotient (IQ). A meta-analysis by Kerr-Wilson, Mackay, Smith, & Pell (2012), covering studies from 1980 to 2009, found preterm delivery to be associated with a 12 point reduction in IQ scores. Therefore, it seems that preterm children, although within the normal range (Johnson, 2007), showed lower IQ scores than their term peers and that improvements in obstetric and neonatal practice have not reduced the impact of preterm delivery on IQ.

Domain-specific measures of neuropsychological abilities have also been studied although it is obvious that some of these cognitive functions may be strongly related to IQ. Executive dysfunctions are commonly evidenced in preterm samples (Mulder, Pitchford, Hagger, & Marlow, 2009); rather than a global deficit, they constitute a single profile of affected and non-affected areas. Executive function abnormalities comprise: (a) anticipation and deployment of attention; (b) impulse control and self-regulation; (c) initiation of activity; (d) working memory; (e) mental flexibility and utilization of feedback; (f) planning ability and organization, and (g) selection of efficient problem-solving strategies. Moreover, very preterm children are at high risk for executive dysfunctions such as working memory deficits and processing speed dysfunctions which may impact academic achievement and inattention and impulsive behavior (Mulder *et al.*, 2009; Mulder, Pitchford, & Marlow, 2010; Mulder, Pitchford, & Marlow, 2011).

Preterm birth also negatively affects visual recognition memory, visuospatial memory and visuospatial perception in childhood (Jongbloed-Pereboom, Janssen, Steenbergen, & Nijhuis-van der Sanden, 2012). As regards the language development in children born preterm without cerebral damage, linguistic abnormalities were found at the end of the preschool years (Guarini *et al.*, 2009) in addition to language (reading comprehension, lexical production and phonological awareness) and literacy problems (reading comprehension, reading and writing) by the age of eight years (Guarini *et al.*, 2010).

In addition to cognitive problems, preterm children are at high risk for behavioral problems (Bhutta *et al.*, 2002). Attention deficit-hyperactivity disorder (ADHD) is the primary abnormal outcome. Specifically, there is a 23% prevalence and nearly threefold increased risk for psychiatric disorders such as ADHD, emotional problems and autism spectrum disorders at 11 years of age in extremely preterm children (Johnson *et al.*, 2007).

Neuropsychological profile of children born preterm with periventricular leukomalacia

Periventricular leukomalacia (PVL) is the prototypical lesion in the encephalopathy of prematurity and the most prevalent type of brain injury affecting cerebral

white matter (WM). It leads to long-lasting cognitive and sensory-motor impairments in this population; 25-50% of very low birth weight infants present cognitive and behavioral deficits and 5-10% present cerebral palsy (Volpe, 2008, 2009).

It has been shown that preterm children who suffered a neonatal periventricular injury display sub-optimal scores compared to full-term children in performance and verbal intelligence quotients at 11 years. Those with severe damage evidenced worse reading and spelling skills, which are both related to working memory deficits (Downie, Frisk, & Jakobson, 2005). High-risk preterm children with poor reading skills may be likely to show executive dysfunctions (Frye, Landry, Swank, & Smith, 2009). Low scores for mental calculation have also been found in adolescents who suffered periventricular brain damage (Pavlova, Sokolov, & Krageloh-Mann, 2009).

Selective deficits in visuospatial and visuomotor integration have been found in preterm children with PVL (Olsen *et al.*, 1998) and are attributed to a malfunctioning of the dorsal stream (Fazzi *et al.*, 2004). Interestingly, Fazzi *et al.* (2009) have pointed out that PVL preterm children at 8 years showed involvement of both dorsal and ventral streams, reflected in deficits mainly in visual-object recognition, visual imagery, visuospatial skills and visual memory.

There are two main possibilities for explaining cognitive deficits after PVL (Kostovic & Judas, 2006). First, diffuse perinatal WM lesions can be so extensive as to involve a variety of motor, sensory and corticocortical pathways (Counsell *et al.*, 2003) and second, the focal component selectively affects a topographically condensed periventricular crossroad of mixed motor, sensory and associative pathways (Judas *et al.*, 2005). Thus, this combination of extensive and more restrictive damage may be linked with the variable cognitive performance evidenced in infants with perinatal brain lesions.

Predictors of long-term cognitive outcome in children/adolescents born preterm

The nature of cognitive development in the preterm infant is influenced by genetic, sociodemographic and congenital factors (Luciana, 2003). Among the main predictors of long-term outcome during childhood and adolescence, brain immaturity, lower birth weight (BW) and gestational age (GA) have been commonly reported (Bhutta *et al.*, 2002). It has been reported that early GA predicts the loss of around 1.5 to 2.5 IQ points per week in very preterm children (Johnson, 2007). In a recent study by the EPIPAGE group (Beaino *et al.*, 2011), the authors concluded that social factors such as low parental socio-economic status and lack of breastfeeding were significant predictors of mild cognitive dysfunctions. Both these factors, as well as biological factors including the presence of cerebral lesions, being small for GA and having a large number of siblings predicted severe cognitive deficiency. In addition, in a sample of extremely low BW

children maternal education was found to be the strongest predictor of long-term neurodevelopmental outcome between 6 and 10-13 years of age, followed by the presence of brain damage (Voss, Jungmann, Wachtendorf, & Neubauer, 2012). Another biological factor such as early motor repertoire has been identified as a good marker for intelligence performance from 7 to 11 years (Bruggink, Van Braeckel, & Bos, 2010). As far as executive functions are concerned, postnatal growth (for spatial span and planning) and parental education (for verbal fluency) were the main predictors of executive functions (Aarnoudse-Moens, Weisglas-Kuperus, Duivenvoorden, Oosterlaan, & van Goudoever, 2013).

A good predictor of IQ in early adolescence is a previous IQ score at an early age (Botting, Powls, Cooke, & Marlow, 1998). However, it is still not clear whether early childhood cognitive performance is predictive of long-term consequences, since numerous cognitive functions emerge later in life and others are in the early stages of development (Anderson & Doyle, 2008). Furthermore, longitudinal studies have not found evidence of catch-up growth in comparison with term peers; rather, worsening outcomes have been reported over time (Johnson, 2007). In contrast, in a longitudinal study comprising a sample of very low BW infants Ment *et al.* (2003) found an improvement in verbal and IQ test scores in middle childhood.

MRI studies of brain morphology: Update and their cognitive correlates in preterm children and adolescents

Sulcal impairments

MRI techniques have been used in preterm subjects to investigate sulcal and gyral abnormalities. In subjects with evident abnormalities such as parenchymal hemorrhages and cystic PVL (Battin *et al.*, 1998), preterm birth seems to affect the staging of normal gyrification; furthermore, when quantitative measures are used, impairments in gyration are seen even in subjects without perinatal complications (Kesler *et al.*, 2006; Gimenez *et al.*, 2006a). Kesler *et al.* (2006) reported that children born prematurely showed an impaired gyrification index in the temporal lobe, affecting reading skills. In an adolescent sample, Gimenez *et al.* (2006a) found a significantly lower maximum depth in the orbitofrontal sulcus, but not in the primary olfactory sulcus.

The cause of the sulcal abnormalities associated with prematurity is still unknown. It has been suggested that the specific location and shape of sulci are determined by the global minimization over the brain of the visco-elastic tensions from WM fibers connecting cortical areas (Van Essen, 1997). This question was addressed by Dubois *et al.* (2008a), who compared 25 preterm subjects with clinically normal MRI and 10 subjects with WM lesions, observing a trend towards higher sulcation indices in the frontal lobe in subjects with WM lesions.

Regional gray and white matter abnormalities

Areas previously associated with GM abnormalities using quantitative volumetric MRI techniques were: temporal lobe bilaterally, parietal cortices (Peterson *et al.*, 2000; Soria-Pastor *et al.*, 2009), frontal and occipital areas (Nosarti *et al.*, 2008), cerebellum (Allin *et al.*, 2001), caudate nucleus (Abernethy, Cooke, & Foulder-Hughes, 2004), thalamus (Nosarti *et al.*, 2008) and hippocampus (Gimenez *et al.*, 2004). Impairment of any of these areas has the potential to disrupt normal neurodevelopment. MRI studies during childhood and adolescence have shown WM reductions in the brainstem (Hargitai *et al.*, 2004), internal capsule (Partridge *et al.*, 2004), subthalamic nuclei, the pons, temporal and frontal regions (Gimenez *et al.*, 2006b) and the corpus callosum (Peterson *et al.*, 2000). These differences in GM and WM volumes appear to be gender-specific, with males presenting greater impairment (Kesler *et al.*, 2008) and directly related to neonatal factors (GA and BW) during childhood (Soria-Pastor *et al.*, 2009).

Although cortical and subcortical regions with reduced GM have been described in preterm samples at school age and in adolescence, few reports have isolated samples of preterm children with PVL in order to investigate cerebral GM at this time of life. Interest in this field is increasing, and a recent review concluded that PVL WM injury is often accompanied by diffuse neuronal/axonal disease affecting cortical and subcortical GM (Volpe, 2009). The first study to provide evidence of GM damage in association with PVL was carried out by Inder *et al.* (1999) in a group of 10 neonates with this condition, highlighting that decrements in total myelinated WM were accompanied by global cortical GM reductions. Although PVL characteristically affects cerebral WM (Kwaja & Volpe, 2008), neuropathological studies have shown that GM lesions are frequent in infants with PVL, indicating that WM injury is often accompanied by GM abnormalities (Pierson *et al.*, 2007). Quantification of neuronal loss in a PVL post-mortem study showed that the structure with the largest GM reduction was the thalamus, which presented a loss of 38% compared with one of 21% in the cerebral cortex (Pierson *et al.*, 2007). Thalamic damage is also a common finding in PVL neuropathological studies (Pierson *et al.*, 2007; Ligam *et al.*, 2009), and in MRI studies (Ricci *et al.*, 2006); its dorsomedial and pulvinar regions have been reported to be the most frequently affected areas (Nagasunder *et al.*, 2011).

Previous quantitative MRI studies in preterm children found that regional GM changes in regions such as sensorimotor, middle temporal and postcentral gyri correlated with IQ scores (Peterson *et al.*, 2000; Soria-Pastor *et al.*, 2009). It has also been reported that frontal and temporal volume decrements correlate with verbal IQ decline in preterm children, while decreases in temporal and occipital cortical regions are associated with a decline in performance IQ (Isaacs *et al.*, 2004). White matter volume has been directly related to performance IQ in preterm adolescents (Soria-Pastor *et al.*, 2008). In PVL children, neurodevelopmental outcome

correlated with the severity of MRI findings (Serdaroglu, Tekgul, Kitis, Serdaroglu, & Gokben, 2004). Moreover, parieto-occipital periventricular lesions of preterm children with PVL lead to long-term disturbances in visual perceptual system functioning (Pavlova, Sokolov, Birbaumer, & Krageloh-Mann, 2006) whereas frontal lesions in the right hemisphere are linked with visual navigation disabilities (Pavlova *et al.*, 2007), and right temporal periventricular lesions are related to the ability for perception and understanding of others' actions (Pavlova, Sokolov, Birbaumer, & Krageloh-Mann, 2008).

Increases and decrements in cortical thickness

Normal development in childhood is accompanied by an increase in CTh (Cortical Thickness) in almost all brain areas, whereas during adolescence there is a progressive cortical thinning (Shaw *et al.*, 2008). Studies of neurodevelopmental disorders have reported both thicker and thinner cortical regions. For instance, cortical thickening has been described in autism (Raznahan *et al.*, 2010) whereas cortical thinning has been found in ADHD (Shaw *et al.*, 2007).

Two previous studies in preterm adolescents measuring CTh have shown cortical thickening and thinning, and a greater impact with lower BW and shorter GA (Martinussen *et al.*, 2005; Nagy Lagercrantz, & Hutton, 2011). Intelligence levels correlated with CTh maturation trajectories in both children and adolescents (Shaw *et al.*, 2006). A recent study performed in a large sample of normal developing brains found positive bilateral correlations between intelligence and CTh localized in frontal, parietal, temporal and occipital lobes in adolescence (Karama *et al.*, 2012). Moreover, positive correlations have been found between IQ and CTh in areas in ventro-lateral frontal, parietal and temporal lobes in very low BW late teenagers (Bjuland, Lohaugen, Martinussen, & Skranes, 2013), and entorhinal cortical thinning has been associated with low IQ and reduced perceptual and executive functions in preterm adolescents (Skranes *et al.*, 2012).

REFERENCES

- Aarnoudse-Moens, C.S., Weisglas-Kuperus, N., Duivenvoorden, H.J., Oosterlaan, J., & van Goudoever, J.B. (2013). Neonatal and parental predictors of executive function in very preterm children. *Acta Paediatrica*, *102*(3):282-286.
- Abernethy, L.J., Cooke, R.W., & Foulder-Hughes, L. (2004). Caudate and hippocampal volumes, intelligence, and motor impairment in 7-year-old children who were born preterm. *Pediatric Research*, *55*(5), 884-893.
- Allin, M., Matsumoto, H., Santhouse, A.M., Nosarti, C., AlAsady, M.H., Stewart, A. L., . . . Murray, R.M. (2001). Cognitive and motor function and the size of the cerebellum in adolescents born very pre-term. *Brain: 124*(Pt 1), 60-66.

- Anderson, P.J., & Doyle, L.W. (2008). Cognitive and educational deficits in children born extremely preterm. *Seminars in Perinatology*, 32(1), 51-58.
- Aylward, G.P. (2002). Cognitive and neuropsychological outcomes: More than IQ scores. *Mental Retardation and Developmental Disabilities Research Reviews*, 8(4), 234-240
- Aylward, G.P. (2005). Neurodevelopmental outcomes of infants born prematurely. *Journal of Developmental and Behavioral Pediatrics : JDBP*, 26(6), 427-440.
- Battin, M.R., Maalouf, E.F., Counsell, S.J., Herlihy, A.H., Rutherford, M.A., Azzopardi, D., & Edwards, A.D. (1998). Magnetic resonance imaging of the brain in very preterm infants: Visualization of the germinal matrix, early myelination, and cortical folding. *Pediatrics*, 101(6), 957-962.
- Beaino, G., Khoshnood, B., Kaminski, M., Marret, S., Pierrat, V., Vieux, R., . . . EPIPAGE Study Group. (2011). Predictors of the risk of cognitive deficiency in very preterm infants: The EPIPAGE prospective cohort. *Acta Paediatrica*, 100(3), 370-378.
- Bhutta, A.T., Cleves, M.A., Casey, P.H., Cradock, M.M., & Anand, K.J. (2002). Cognitive and behavioral outcomes of school-aged children who were born preterm: A meta-analysis. *JAMA : The Journal of the American Medical Association*, 288(6), 728-737.
- Bjuland, K.J., Lohaugen, G.C., Martinussen, M., & Skranes, J. (2013). Cortical thickness and cognition in very-low-birth-weight late teenagers. *Early Human Development* ;89 (6):371-380
- Blondel, B., & Kaminski, M. (2002). Trends in the occurrence, determinants, and consequences of multiple births. *Seminars in Perinatology*, 26(4), 239-249.
- Botting, N., Powls, A., Cooke, R.W., & Marlow, N. (1998). Cognitive and educational outcome of very-low-birthweight children in early adolescence. *Developmental Medicine and Child Neurology*, 40(10), 652-660.
- Bruggink, J.L., Van Braeckel, K.N., & Bos, A.F. (2010). The early motor repertoire of children born preterm is associated with intelligence at school age. *Pediatrics*, 125(6), e1356-1363.
- Counsell, S.J., Allsop, J.M., Harrison, M.C., Larkman, D.J., Kennea, N.L., Kapellou, O., . . . Rutherford, M.A. (2003). Diffusion-weighted imaging of the brain in preterm infants with focal and diffuse white matter abnormality. *Pediatrics*, 112(1 Pt 1), 1-7.
- Downie, A.L., Frisk, V., & Jakobson, L.S. (2005). The impact of periventricular brain injury on reading and spelling abilities in the late elementary and adolescent years. *Child Neuropsychology: A Journal on Normal and Abnormal Development in Childhood and Adolescence*, 11(6), 479-495.
- Dubois, J., Benders, M., Cachia, A., Lazeyras, F., Ha-Vinh Leuchter, R., Sizonenko, S. V., . . . Huppi, P. S. (2008a). Mapping the early cortical folding process in the preterm newborn brain. *Cerebral Cortex*, 18(6), 1444-1454.
- Fazzi, E., Bova, S.M., Uggetti, C., Signorini, S.G., Bianchi, P.E., Maraucci, I., . . . Lanzi, G. (2004). Visual-perceptual impairment in children with periventricular leukomalacia. *Brain & Development*, 26(8), 506-512.
- Fazzi, E., Bova, S., Giovenzana, A., Signorini, S., Uggetti, C., & Bianchi, P. (2009). Cognitive visual dysfunctions in preterm children with periventricular leukomalacia. *Developmental Medicine and Child Neurology*, 51 (12),974-981.
- Frye, R. E., Landry, S. H., Swank, P. R., & Smith, K. E. (2009). Executive dysfunction in poor readers born prematurely at high risk. *Developmental Neuropsychology*, 34(3), 254-271.
- Gimenez, M., Junque, C., Narberhaus, A., Caldu, X., Salgado-Pineda, P., Bargallo, N., . . . Botet, F. (2004). Hippocampal gray matter reduction associates with memory deficits in adolescents with history of prematurity. *Neuroimage*, 23(3), 869-877.
- Gimenez, M., Junque, C., Vendrell, P., Narberhaus, A., Bargallo, N., Botet, F., & Mercader, J. M. (2006a). Abnormal orbitofrontal development due to prematurity. *Neurology*, 67(10), 1818-1822.
- Gimenez M., Junque C., Narberhaus A., Bargallo N., Botet F., & Mercader J.M. (2006b) White matter volume and concentration reductions in adolescents with history of very preterm birth: a voxel-based morphometry study. *Neuroimage*, 32(4):1485-1498.

- Guarini, A., Sansavini, A., Fabbri, C., Alessandrini, R., Faldella, G., & Karmiloff-Smith, A. (2009). Reconsidering the impact of preterm birth on language outcome. *Early Human Development*, 85(10), 639-645.
- Guarini, A., Sansavini, A., Fabbri, C., Savini, S., Alessandrini, R., Faldella, G., & Karmiloff-Smith, A. (2010). Long-term effects of preterm birth on language and literacy at eight years. *Journal of Child Language*, 37(4), 865-885.
- Hargitai, B., Szabo, V., Cziniel, M., Hajdu, J., Papp, Z., Szende, B., & Sergi, C. (2004). Human brain of preterm infants after hypoxic-ischaemic injuries: No evidence of a substantial role for apoptosis by using a fine-tuned ultrasound-guided neuropathological analysis. *Brain & Development*, 26(1), 30-36.
- Inder, T.E., Huppi, P.S., Warfield, S., Kikinis, R., Zientara, G.P., Barnes, P.D., . . . Volpe, J.J. (1999). Periventricular white matter injury in the premature infant is followed by reduced cerebral cortical gray matter volume at term. *Annals of Neurology*, 46(5), 755-760.
- Isaacs, E.B., Edmonds, C.J., Chong, W.K., Lucas, A., Morley, R., & Gadian, D.G. (2004). Brain morphometry and IQ measurements in preterm children. *Brain*, 127(Pt 12), 2595-2607.
- Johnson, S. (2007). Cognitive and behavioural outcomes following very preterm birth. *Seminars in Fetal & Neonatal Medicine*, 12(5), 363-373.
- Jongbloed-Pereboom, M., Janssen, A. J., Steenbergen, B., & Nijhuis-van der Sanden, M.W. (2012). Motor learning and working memory in children born preterm: A systematic review. *Neuroscience and Biobehavioral Reviews*, 36(4), 1314-1330.
- Judas, M., Rados, M., Jovanov-Milosevic, N., Hrabac, P., Stern-Padovan, R., & Kostovic, I. (2005). Structural, immunocytochemical, and mr imaging properties of periventricular crossroads of growing cortical pathways in preterm infants. *AJNR.American Journal of Neuroradiology*, 26(10), 2671-2684.
- Karama, S., Colom, R., Johnson, W., Deary, I.J., Haier, R., Waber, D.P., . . . Evans, A.C., & Brain Development Cooperative Group. (2011). Cortical thickness correlates of specific cognitive performance accounted for by the general factor of intelligence in healthy children aged 6 to 18. *Neuroimage*, 55(4), 1443-2453.
- Khwaja, O., & Volpe, J.J. (2008). Pathogenesis of cerebral white matter injury of prematurity. *Archives of Disease in Childhood.Fetal and Neonatal Edition*, 93(2), 153-161.
- Kerr-Wilson, C.O., Mackay, D.F., Smith, G.C., & Pell, J.P. (2012). Meta-analysis of the association between preterm delivery and intelligence. *Journal of Public Health* 34(2), 209-216.
- Kesler, S.R., Vohr, B., Schneider, K.C., Katz, K.H., Makuch, R.W., Reiss, A.L., & Ment, L.R. (2006). Increased temporal lobe gyriification in preterm children. *Neuropsychologia*, 44(3), 445-453.
- Kesler, S.R., Reiss, A.L., Vohr, B., Watson, C., Schneider, K.C., Katz, K.H., . . . Ment, L.R. (2008). Brain volume reductions within multiple cognitive systems in male preterm children at age twelve. *The Journal of Pediatrics*, 152(4), 513-520.
- Kostovic, I., & Judas, M. (2006). Prolonged coexistence of transient and permanent circuitry elements in the developing cerebral cortex of fetuses and preterm infants. *Developmental Medicine and Child Neurology*, 48(5), 388-393.
- Ligam, P., Haynes, R.L., Folkerth, R.D., Liu, L., Yang, M., Volpe, J.J., & Kinney, H C. (2009). Thalamic damage in periventricular leukomalacia: Novel pathologic observations relevant to cognitive deficits in survivors of prematurity. *Pediatric Research*, 65(5), 524-529.
- Luciana, M. (2003). Cognitive development in children born preterm: Implications for theories of brain plasticity following early injury. *Development and Psychopathology*, 15(4), 1017-1047.
- Marlow, N. (2004). Neurocognitive outcome after very preterm birth. *Archives of Disease in Childhood.Fetal and Neonatal Edition*, 89(3), 224-228.
- Martinussen, M., Fischl, B., Larsson, H.B., Skranes, J., Kulseng, S., Vangberg, T.R., . . . Dale, A.M. (2005). Cerebral cortex thickness in 15-year-old adolescents with low birth weight measured by an automated MRI-based method. *Brain*: 128, 2588-2596.

- Ment, L.R., Vohr, B., Allan, W., Katz, K.H., Schneider, K.C., Westerveld, M., . . . Makuch, R.W. (2003). Change in cognitive function over time in very low-birth-weight infants. *JAMA : The Journal of the American Medical Association*, 28;289(6):705-711.
- Mulder, H., Pitchford, N.J., Hagger, M.S., & Marlow, N. (2009). Development of executive function and attention in preterm children: A systematic review. *Developmental Neuropsychology*, 34(4), 393-421
- Mulder, H., Pitchford, N.J., & Marlow, N. (2010). Processing speed and working memory underlie academic attainment in very preterm children. *Archives of Disease in Childhood.Fetal and Neonatal Edition*, 95(4), F267-72.
- Mulder, H., Pitchford, N. J., & Marlow, N. (2011). Processing speed mediates executive function difficulties in very preterm children in middle childhood. *Journal of the International Neuropsychological Society : JINS*, 1-10.
- Nagasunder, A.C., Kinney, H.C., Bluml, S., Tavare, C.J., Rosser, T., Gilles, F.H., . . . Panigrahy, A. (2011). Abnormal microstructure of the atrophic thalamus in preterm survivors with periventricular leukomalacia. *AJNR.American Journal of Neuroradiology*, 32(1), 185-191.
- Nagy, Z., Lagercrantz, H., & Hutton, C. (2011). Effects of preterm birth on cortical thickness measured in adolescence. *Cerebral Cortex*, 21(2), 300-306.
- Nosarti, C., Giouroukou, E., Healy, E., Rifkin, L., Walshe, M., Reichenberg, A., . . . Murray, R. M. (2008). Grey and white matter distribution in very preterm adolescents mediates neurodevelopmental outcome. *Brain*, 131(Pt 1), 205-217.
- Olsen, P., Vainionpaa, L., Paakko, E., Korkman, M., Pyhtinen, J., & Jarvelin, M.R. (1998). Psychological findings in preterm children related to neurologic status and magnetic resonance imaging. *Pediatrics*, 102(2), 329-336.
- Partridge, S.C., Mukherjee, P., Henry, R.G., Miller, S.P., Berman, J.I., Jin, H., . . . Vigneron, D.B. (2004). Diffusion tensor imaging: Serial quantitation of white matter tract maturity in premature newborns. *Neuroimage*, 22(3), 1302-1314.
- Pavlova, M., Sokolov, A.N., Birbaumer, N., & Krageloh-Mann, I. (2006). Biological motion processing in adolescents with early periventricular brain damage. *Neuropsychologia*, 44(4), 586-593.
- Pavlova, M., Sokolov, A.N., Birbaumer, N., & Krageloh-Mann, I. (2008). Perception and understanding of others' actions and brain connectivity. *Journal of Cognitive Neuroscience*, 20(3), 494-504.
- Pavlova, M., Sokolov, A. N., & Krageloh-Mann, I. (2009). Arithmetic and brain connectivity: Mental calculation in adolescents with periventricular lesions. *Neuropsychologia*, 47(2), 439-445.
- Peterson, B. S., Vohr, B., Staib, L. H., Cannistraci, C. J., Dolberg, A., Schneider, K. C., . . . Ment, L. R. (2000). Regional brain volume abnormalities and long-term cognitive outcome in preterm infants. *JAMA : The Journal of the American Medical Association*, 284(15), 1939-1947.
- Pierson, C.R., Folkerth, R.D., Billiards, S.S., Trachtenberg, F.L., Drinkwater, M.E., Volpe, J.J., & Kinney, H.C. (2007). Gray matter injury associated with periventricular leukomalacia in the premature infant. *Acta Neuropathologica*, 114(6), 619-631.
- Raznahan, A., Toro, R., Daly, E., Robertson, D., Murphy, C., Deeley, Q., . . . Murphy, D.G. (2010). Cortical anatomy in autism spectrum disorder: An in vivo MRI study on the effect of age. *Cerebral Cortex*, 20(6), 1332-1340.
- Ricci, D., Anker, S., Cowan, F., Pane, M., Gallini, F., Luciano, R., . . . Mercuri, E. (2006). Thalamic atrophy in infants with PVL and cerebral visual impairment. *Early Human Development*, 82(9), 591-595.
- Serdaroglu, G., Tekgul, H., Kitis, O., Serdaroglu, E., & Gokben, S. (2004). Correlative value of magnetic resonance imaging for neurodevelopmental outcome in periventricular leukomalacia. *Developmental Medicine and Child Neurology*, 46(11), 733-739.

- Shaw, P., Greenstein, D., Lerch, J., Clasen, L., Lenroot, R., Gogtay, N., . . . Giedd, J. (2006). Intellectual ability and cortical development in children and adolescents. *Nature*, *440* (7084), 676-679.
- Shaw, P., Eckstrand, K., Sharp, W., Blumenthal, J., Lerch, J.P., Greenstein, D., . . . Rapoport, J.L. (2007). Attention-deficit/hyperactivity disorder is characterized by a delay in cortical maturation. *Proceedings of the National Academy of Sciences of the United States of America*, *104*(49), 19649-19654.
- Shaw, P., Kabani, N.J., Lerch, J.P., Eckstrand, K., Lenroot, R., Gogtay, N., . . . Wise, S.P. (2008). Neurodevelopmental trajectories of the human cerebral cortex. *The Journal of Neuroscience*, *28*(14), 3586-3594.
- Simmons, L.E., Rubens, C.E., Darmstadt, G.L., & Gravett, M.G. (2010). Preventing preterm birth and neonatal mortality: Exploring the epidemiology, causes, and interventions. *Seminars in Perinatology*, *34*(6), 408-415.
- Skranes, J., Lohaugen, G.C., Evensen, K.A., Indredavik, M.S., Haraldseth, O., Dale, A.M., . . . Martinussen, M. (2012). Entorhinal cortical thinning affects perceptual and cognitive functions in adolescents born preterm with very low birth weight (VLBW). *Early Human Development*, *88*(2), 103-109.
- Soria-Pastor, S., Gimenez, M., Narberhaus, A., Falcon, C., Botet, F., Bargallo, N., . . . Junque, C. (2008). Patterns of cerebral white matter damage and cognitive impairment in adolescents born very preterm. *International Journal of Developmental Neuroscience*, *26*(7), 647-654.
- Soria-Pastor, S., Padilla, N., Zubiaurre-Elorza, L., Ibarretxe-Bilbao, N., Botet, F., Costas-Moragas, C., . . . Junque, C. (2009). Decreased regional brain volume and cognitive impairment in preterm children at low risk. *Pediatrics*, *124*(6), e1161-70.
- Van Essen, D.C. (1997). A tension-based theory of morphogenesis and compact wiring in the central nervous system. *Nature*, *385*(6614), 313-318.
- Volpe, J.J. *Neurology of the Newborn* (4th edition). (2008). Philadelphia, WB Saunders.
- Volpe, J.J. (2009). Brain injury in premature infants: A complex amalgam of destructive and developmental disturbances. *Lancet Neurology*, *8*(1), 110-124.
- Voss, W., Jungmann, T., Wachtendorf, M., & Neubauer, A.P. (2012). Long-term cognitive outcomes of extremely low birth weight infants: The influence of the maternal educational background. *Acta Paediatrica*, *101*(6), 569-573.
- World Health Organization. (1977). WHO: Recommended definitions, terminology and format for statistical tables related to the perinatal period and use of a new certificate for cause of perinatal deaths, modifications recommended by FIGO as amended october 14, 1976. *Acta Obstetrica Et Gynecologica Scandinavica*, *556*(3), 247-253.