Collect. Math. **59**, 3 (2008), 321–324

© 2008 Universitat de Barcelona

A local form for the automorphisms of the spectral unit ball

PASCAL J. THOMAS

Institut de Mathématiques de Toulouse, UMR CNRS 5219

Université Paul Sabatier, 118 Route de Narbonne

F-31062 Toulouse Cedex, France

E-mail: pthomas@cict.fr

Received January 21, 2008. Revised February 13, 2008.

ABSTRACT

If F is an automorphism of Ω_n , the n^2 -dimensional spectral unit ball, we show that, in a neighborhood of any cyclic matrix of Ω_n , the map F can be written as conjugation by a holomorphically varying non singular matrix. This provides a shorter proof of a theorem of J. Rostand, with a slightly stronger result.

1. Background

Let $\mathcal{M}_n$ be the set of all $n \times n$ complex matrices. For $A \in \mathcal{M}_n$ denote by $sp(A)$ the spectrum of A . The spectral ball Ω_n is the set

$$\Omega_n := \{A \in \mathcal{M}_n : \forall \lambda \in sp(A), |\lambda| < 1\}.$$

Let F be an automorphism of Ω_n , that is to say, a biholomorphic map of the spectral ball into itself. Ransford and White [6] proved that, by composing with a natural lifting of a Möbius map of the disk, one could reduce oneself to the case where $F(0) = 0$, and that in that case the linear map $F'(0)$ was a linear automorphism of Ω_n , so that by composing with its inverse, one is reduced to the case $F(0) = 0$, $F'(0) = I$ (the identity map). We then say that the automorphism is *normalized*. Ransford and White [6] proved that such automorphisms preserve the spectrum of matrices.

We say that two matrices X, Y are *conjugate* if there exists $Q \in \mathcal{M}_n^{-1}$ such that $X = Q^{-1}YQ$.

This paper was made possible in part by a grant from the French Ministry of Foreign Affairs in the framework of the ECO Net programme, file 10291 SL, coordinated by Ahmed Zeriahi.

Keywords: Holomorphic automorphisms, spectral ball, non-derogatory matrices, cyclic matrices.

MSC2000: 32H02, 32A07, 15A21.

Baribeau and Ransford [1] (see also [2] for a more elementary proof) proved that every spectrum-preserving $\mathcal{C}^1$ -diffeomorphism of an open subset of $\mathcal{M}_n$, and thus every normalized automorphism of the spectral ball is a pointwise conjugation:

$$F(X) = Q(X)^{-1}XQ(X). \quad (1)$$

Rostand's contribution [7] was to show that $Q(X)$ could be chosen locally holomorphically in a neighborhood of every X admitting n distinct eigenvalues.

We will give a short proof of a slightly stronger result: the exceptional set of matrices where the local holomorphic choice cannot be guaranteed will be of complex codimension 2 instead of 1.

The motivation for this result was a conjecture formulated in [6] about the automorphisms of the spectral ball, which reduces to asking whether any normalized automorphisms can be written in the form (1), where Q would be globally holomorphic on Ω_n , and depend only on the conjugacy class of X . Notice that a recent result of Zwonek [8] shows that any proper map of the spectral ball to itself is actually an automorphism of it, so that the proof of the Ransford-White conjecture would yield a description of all the proper maps of the spectral ball into itself.

I wish to thank Nikolai Nikolov, who told me about this circle of ideas. Without the fruitful discussions I had with him on this topic, this paper wouldn't have been written.

2. Statement

DEFINITION 1 We say that a matrix M is *cyclic* (or *non-derogatory*) if there exists a cyclic vector for M , i.e. $v \in \mathbb{C}^n$ such that $(v, Mv, \dots, M^{k-1}v, \dots)$ spans $\mathbb{C}^n$, which is equivalent to the fact that $(v, Mv, \dots, M^{n-1}v)$ is a basis of $\mathbb{C}^n$.

Many equivalent definitions of this notion can be found, for instance in [3, 4], or [5, Proposition 3]. We point one out: M is cyclic if and only if for any $\lambda \in \mathbb{C}$, $\dim \text{Ker}(M - \lambda I_n) \leq 1$. In particular, any matrix with n distinct eigenvalues is cyclic, and for any given spectrum $\lambda_1, \dots, \lambda_n$, the set of non-cyclic matrices with that spectrum is the algebraic set

$$\{M : \exists j : \dim \text{Ker}(M - \lambda_j I_n) \geq 2\}.$$

Hence the set of non-cyclic matrices is of codimension 1 in the set of matrices which admit at least one multiple eigenvalue, itself of codimension 1 in $\mathcal{M}_n$.

Theorem 2

Let F be a spectrum-preserving holomorphic map of Ω_n . Let $X_0 \in \Omega_n$ be a cyclic matrix. Then there exists a neighborhood $\mathcal{V}_{X_0}$ of X_0 and a map Q holomorphic from $\mathcal{V}_{X_0}$ to $\mathcal{M}_n^{-1}$ such that for any $X \in \mathcal{V}_{X_0}$, $F(X) = Q(X)^{-1}XQ(X)$.

3. Proof

We fix some notation. For $A \in \mathcal{M}_n$, let

$$\sigma_j(A) := \sigma_j(\lambda_1, \dots, \lambda_n) := (-1)^j \sum_{1 \leq k_1 < \dots < k_j \leq n} \lambda_{k_1} \dots \lambda_{k_j}$$

and $\lambda_1, \dots, \lambda_n$ are the (possibly equal) eigenvalues of A . Those are polynomials in the coefficients of A . Put $\sigma := (\sigma_1, \dots, \sigma_n) : \mathcal{M}_n \rightarrow \mathbb{C}^n$.

Conversely, given $a := (a_1, \dots, a_n) \in \mathbb{C}^n$, the associated *companion matrix* $\mathcal{C}_a$ is

$$\begin{pmatrix} 0 & & & -a_n \\ 1 & 0 & & \vdots \\ & 1 & \ddots & \vdots \\ & & \ddots & 0 & -a_2 \\ & & & 1 & -a_1 \end{pmatrix}.$$

The companion matrix associated to a matrix A is $\mathcal{C}_{\sigma(A)}$. They have the same characteristic polynomial, or equivalently $\sigma(\mathcal{C}_{\sigma(A)}) = \sigma(A)$.

Now given a matrix X_0 as in the Theorem, and a vector v_0 cyclic for X_0 , let

$$\mathcal{U}_{X_0} := \{M \in \mathcal{M}_n : \det(v_0, Mv_0, \dots, M^{n-1}v_0) \neq 0\}.$$

This is a neighborhood of X_0 . Let $P_{v_0}(M)$ be the matrix with columns $(v_0, Mv_0, \dots, M^{n-1}v_0)$; this depends polynomially on the entries of M , and is invertible. One can see that for $X \in \mathcal{U}_{X_0}$,

$$P_{v_0}(X)^{-1}XP_{v_0}(X) = \mathcal{C}_{\sigma(X)}$$

(the $n-1$ st columns coincide, and they have the same characteristic polynomial).

By the Baribeau-Ransford theorem [1] $F(X_0)$ is conjugate to X_0 , therefore cyclic. So there is a vector w_0 , and a neighborhood $\mathcal{U}_{F(X_0)}$ where the relation $P_{w_0}(Y)^{-1}YP_{w_0}(Y) = \mathcal{C}_{\sigma(Y)}$ holds. Take $\mathcal{V}_{X_0} \subset \mathcal{U}_{X_0}$ small enough so that $F(\mathcal{V}_{X_0}) \subset \mathcal{U}_{F(X_0)}$. For any $X \in \mathcal{V}_{X_0}$, using the fact that F is spectrum-preserving,

$$\begin{aligned} F(X) &= P_{w_0}(F(X))\mathcal{C}_{\sigma(F(X))}P_{w_0}(F(X))^{-1} \\ &= P_{w_0}(F(X))\mathcal{C}_{\sigma(X)}P_{w_0}(F(X))^{-1} \\ &= P_{w_0}(F(X))P_{v_0}(X)^{-1}XP_{v_0}(X)P_{w_0}(F(X))^{-1}, \end{aligned}$$

so that we have the theorem with $Q(X) = P_{v_0}(X)P_{w_0}(F(X))^{-1}$. □

Notice that there is no hope to make a global holomorphic choice of v_0 on the whole open set of cyclic matrices. Indeed, since the complement of this is of codimension 2, we could then extend it to the whole of Ω_n by Hartog's phenomenon, but it would mean that all matrices are cyclic, which is obviously false.

References

1. L. Baribeau and T.J. Ransford, Non-linear spectrum-preserving maps, *Bull. London Math. Soc.* **32** (2000), 8–14.
2. L. Baribeau and S. Roy, Caractérisation spectrale de la forme de Jordan, *Linear Algebra Appl.* **320** (2000), 183–191.
3. R.A. Horn and C.R. Johnson, *Matrix Analysis*, Cambridge University Press, Cambridge, 1985.
4. R.A. Horn and C.R. Johnson, *Topics in Matrix Analysis*, Cambridge University Press, Cambridge, 1991.
5. N. Nikolov, P.J. Thomas, and W. Zwonek, Discontinuity of the Lempert function and the Kobayashi–Royden metric of the spectral ball, (preprint, 2007), (arXiv:math.CV/0704.2470).
6. T.J. Ransford and M.C. White, Holomorphic self-maps of the spectral unit ball, *Bull. London Math. Soc.* **23** (1991), 256–262.
7. J. Rostand, On the automorphisms of the spectral unit ball, *Studia Math.* **155** (2003), 207–230.
8. W. Zwonek, Proper holomorphic mappings of the spectral unit ball, (preprint, 2007), (arXiv:math.CV/0704.0614).